


CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

CATTLEMAX™

POUR-ON FOR BEEF AND DAIRY CATTLE

ACTIVE CONSTITUENT: 5 g/L IVERMECTIN



For the treatment and control of Ivermectin-sensitive internal and external parasites of beef and dairy cattle.

NIL WITHHOLDING PERIOD FOR MILK



Dalgety
Animal Health

20 L

dalgetyah.nutrien.com.au

101830_0620

CATTLEMAX™

POUR-ON FOR BEEF AND DAIRY CATTLE

For the treatment and control of Ivermectin sensitive:

Gastrointestinal nematodes:

- Adult and immature (L4): Small Brown Stomach Worm (including inhibited stages) (*Ostertagia ostertagi*), Barber's Pole Worm (*Haemonchus placei*), Stomach Hair Worm (*Trichostrongylus axei*), Intestinal Hair Worm (*T. colubriformis*), Small Intestinal Worm (*Cooperia* spp.), Nodule Worm (*Oesophagostomum radiatum*); and
- Adult stages of *Cooperia oncophora*, *C. punctata*, *Oesophagostomum venulosum*, Whip Worm (*Trichuris* spp.), Intestinal Threadworm (*Strongyloides papillosus*); and
- Immature stages (L4) of Thin Necked Intestinal Worm (*Nematodirus* spp.).

Lungworms (adult and immature) *Dictyocaulus viviparus*.

Eyeworm *Thelazia* spp. (adult stages).

Sucking and Biting Lice: *Linognathus vituli*, *Haematopinus eurysternus*, *Bovicola bovis*, *Solenopotes capillatus*.

Mites: *Chorioptes bovis*, *Sarcoptes scabiei var bovis*.

Buffalo Flies *Haematobia irritans exigua*

Cattle Ticks: (*Boophilus microplus*) including organophosphate, synthetic pyrethroid- and amidine-resistant strains. Maximum tick efficacy is achieved 4-5 days after treatment. Engorged female ticks that drop from cattle in the initial days after treatment may lay viable eggs. This should be taken into account when planning a strategic tick control program. CATTLEMAX should be alternated with conventional dips or pour-ons when the programme requires repeated treatments.

PERSISTENT ACTIVITY

CATTLEMAX applied at the recommended dose of 1 mL per 10 kg of liveweight controls infections of Stomach Hair Worm (*Trichostrongylus axei*) and Small Intestinal Worm (*Cooperia* spp.) for up to 14 days after treatment; Small Brown Stomach Worm (*Ostertagia* spp.) and Nodule Worm (*Oesophagostomum radiatum*) for up to 21 days after treatment; Lungworm (*Dictyocaulus viviparus*) for up to 28 days after treatment; Buffalo Fly (*Haematobia irritans exigua*) for up to 21 days after treatment.

For best results CATTLEMAX should be part of a total parasite control programme including internal and external parasites. Consult your veterinarian for a complete parasite control programme.

DIRECTIONS FOR USE

RESTRAINT: DO NOT USE in pregnant cows less than 7 days before calving.

DOSAGE

This product is formulated only for external application to cattle; do not use in other species. The formulation should be applied along the topline in a narrow strip extending from the withers to the tailhead. The dose is 1mL for each 10kg of liveweight.

Dose the mob according to the heaviest animal by liveweight in each group (cows, bulls, calves, heifers etc.).

When there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid overdosing. DO NOT underdose. Cattle should be weighed with either scales or with a weightband.

Liveweight (kg)	Dose (mL)	This pack treats (head)	Liveweight (kg)	Dose (mL)	This pack treats (head)
< 100	1mL/10kg	>2000	351 – 400	40	500
101 – 150	15	1333	401 – 450	45	444
151 – 200	20	1000	451 – 500	50	400
201 – 250	25	800	501 – 550	55	363
251 – 300	30	666	551 – 600	60	333
301 – 350	35	571	601 – 650	65	307

Heavier animals (for example, mature bulls) should receive an additional 5mL for each 50kg over 650kg. Treatment should not be repeated in less than 42 days.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 28 days before slaughter for human consumption.

MILK: NIL.

Bobby calves: DO NOT SLAUGHTER for 4 days bobby calves born to cows treated during pregnancy.

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT SLAUGHTER for export for 28 days after treatment.

SAFETY TO CATTLE: Field studies have demonstrated a wide safety margin. The topically applied formulation is expected to be well tolerated by breeding animals.

GENERAL INSTRUCTIONS

Administration

Efficacy of CATTLEMAX is not adversely affected when the hide is wet or if rainfall occurs shortly after treatment. However, it is not considered good agricultural practice to treat animals if they are wet or rain is likely.

PRECAUTIONS

Use only in well-ventilated areas or outdoors. Close container when not in use.

Cloudiness may result when CATTLEMAX is stored at temperatures below 0°C. Allowing to warm at room temperature will restore the normal appearance without affecting efficacy.

The antiparasitic activity of ivermectin will be impaired if the formulation is applied to areas of skin with mange scabs or lesions, or with dermatoses or adherent materials, eg. caked mud or manure.

This product is for application to the skin surface only. DO NOT give orally or parenterally.

CATTLEMAX is not recommended for use in species other than cattle

NOTE TO USER: The colour of CATTLEMAX may fade when exposed to light. This loss of colour does not reflect loss of potency of ivermectin. However, prolonged exposure (ie. weeks) to light can result in a gradual decline of ivermectin potency in the formulation.

DANGER

GIVES OFF HIGHLY FLAMMABLE VAPOUR. KEEP AWAY FROM HEAT, SPARKS, OPEN FLAME OR OTHER SOURCES OF IGNITION. KEEP CLOSED WHEN NOT IN USE.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT: Ivermectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container. When used according to label directions CATTLEMAX is not likely to have an unintended effect that is harmful to the environment or have an overall adverse effect on dung beetle or earthworm populations. Increased mortality and impaired development of dung beetle larvae may be expected, but for a limited period after treatment. There may also be short-term indirect effect on the rate of earthworm colonisation of dung pats.

SAFETY DIRECTIONS: Poisonous if swallowed. May irritate the eyes. Avoid contact with eyes. Wash hands after use.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26. For further information, refer to the Material Information Data Sheet.

DISPOSAL: Triple or (preferably) pressure rinse container. DO NOT dispose of undiluted chemicals on site. Break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots. Empty containers and product should not be burnt.

STORAGE: Store below 30°C (room temperature). Store bottle to protect from light.

CONDITIONS OF SALE: The use of this product is beyond the control of Nutrien Ag Solutions Limited. Any provisions or rights under the Australian Consumer Law which cannot be excluded by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies expressed or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application of this product are excluded. Nutrien Ag Solutions Limited does not accept any liability (including consequential loss and/or negligence) for any loss or damage connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: May be fatal if swallowed and enters airways; IF SWALLOWED immediately call a POISON CENTER/ doctor; Do NOT induce vomiting; Wear protective gloves/protective clothing/eye protection/face protection; Combustible liquid; No smoking; Keep away from heat, hot surfaces, sparks, open flames and other ignition sources; In case of fire: Use for extinction: CO₂, powder or water spray; Store locked up; Store in a well-ventilated place. Keep cool; Dispose of contents/container in accordance with local/regional/national regulations.

© Registered trade mark of Nutrien Ag Solutions Limited



UN 1219
ISOPROPANOL
(ISOPROPYL ALCOHOL),
FLAMMABLE, LIQUID, N.O.S.
HAZCHEM: 2[Y]E PG: II

IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE	FOR SPECIALIST ADVICE IN AN EMERGENCY DIAL 1800 033 111 24 HOURS AUSTRALIA WIDE
---	--



Nutrien Ag Solutions Limited
Level 5, Building A, 26 Talavera Road,
Macquarie Park, NSW, 2113 • Tel: (02) 9889 5400
Product Support Tel: 1800 44 88 92
APVMA Approval No.: 61414/0808 101840_0620

B:

Expiry: