




160 mm

ROLL DIRECTION FIG 6.

Check:	□ Changes & re-submit Comments:		CHLY AS FINAL RESPONSIBI  Approved, OK to proceed  Name:  Sign:  Date:		Whilst all care has been taken to present this proof and it's details correctly, it is the customers responsibility to ensure that all details meet their requirements. Your signature indicates your acceptance that all is correct. If not, please return proof unsigned and with corrections clearly marked.	
N Pacific L	ahels	F/O No 34627	ART VERSION 03	DATE 18.6.20	020	Pre Press: D. Harper
A.B.N 93 000 881 425	25 ng NSW 2259	BLACK	MAGENTA	PMS 348		
Ph: (02) 4359 7000 Fax: (0		CYAN	YELLOW			
Email: sales@pacificla	bels.com.au	Die-line only (not pr	inted)	PLEASE NOTE: C	olours may vary up to	10% from printed proof.



70 mm

ROLL DIRECTION FIG 6.

PLEASE CHE	CK THIS PRO	OOF THOROUGH	ILY AS FINAL	RESPONS	<b>IBILITY FO</b>	R ACCURACY RI	ESTS WITH YOU.	
Check:	☐ Changes & re-submit		☐ Approved, OK to proceed		<b>d</b> Whilst all	Whilst all care has been taken to present this proof and		
☐ 1. Content			Name:		••	it's details correctly, it is the customers responsibility to		
2. Colours			Sign:		signature	ensure that all details meet their requirements. Your signature indicates your acceptance that all is correct If not, please return proof unsigned and with corrections clearly marked.		
☐ 3. Barcode (Numbers)			Date:					
N Pacific	Labels	F/O No <b>34626F</b>	ART VERSION	04 DATE	17.6.2020		Pre Press: D.Harper	
<b>A.B.N 93 000 881 42</b> 2 Lucca Road, Wyor		BLACK						
Ph: (02) 4359 7000 Fax								
Email: sales@pacificlabels.com.au		Die-line only (not prin	ited)	_ PI FASE	NOTE: Colou	rs may vary up to 10%	6 from printed proof	

00 mm

READ THE ENCLOSED LEAFLET BEFORE USING THIS PRODUCT MILK: DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

TRADE ADVICE - EXPORT SLAUGHTER INTERVAL DIRECTIONS FOR USE DIRECTUMS FOR USE
RESTRAINT:
DO NOT USE in ewes, which are producing or may in
the future produce milk that may be used or processed
for human consumption.
RETREATMENT INTERVIAL Must not be retreated for at
least 10 least five administration. TRADE ADVICE - EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 133 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Landmark on 1300 255 347 or the APVMA website (www.apvma.gov.au/residues).

SAFEIY DIRECTIONS: Poisonous if swallowed. May intitute the nose and throot. Will intitute eyes and stin. Avaid carbot, with pues and distin. Avaid carbot with pues and distin. least 91 days after administration.

CONTRAINDICATION: MOXIMAX MOXIDECTIN LONG ACTING INJECTION FOR USE IN SHEEP is not recommended for use in goats as safety and efficacy have not been evaluated.
DOSAGE AND ADMINISTRATION: Use the contents within irritute the nose and throat. Will irritate eyes and skin. Avoid contact with eyes and skin. Do not inhale. Wash hands after use.

HRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre, Phone Australia 131126.

Additional safety information:

Avoid self-injection. If self-injection occurs, seek medical advice os soon as possible. Additional information is in the Safety Data Sheet (SDS).

ENVIRONMENTAL PROTECTION

Very travic to aquants life. Do not contaminate dams. 6 months of first broaching of the vial. Discard the 6 months of hist producing of the viol, Discard the unused portion.

DOSE RATE: 1 mg moxidectin/kg liveweight (1 mL / 20 kg liveweight)

MOXIMAX MOXIDECTIN LONG ACTING INJECTION FOR SHEEP must be administered by subcutaneous (under the skin) injection at the base of the ear or high on the neck behind the ear. The product is ready to use. Administer the dose according to the dosage table using the special Moximax Moxidectin LA sheep injector gun. Do not use Very toxic to aquatic life. Do not contaminate dams, rivers, streams or other waterways with the any other injector. Weight range | Dose volume | Weight range | Dose volume chemical or used container. DISPOSAL: Bury destroyed empty containers at a depth TOP COART COMP QUESTIONED PRINTY CONTINUES OF ORDER

STORE TO THE COART \*DO NOT USE IN LAMBS LESS THAN 20 KG temperature). Protect from light. Nutrien Ag Solutions Limited Level 5, Building A, 26 Talavera Road, When MOXIMAX MOXIDECTIN LONG ACTING INJECTION FOR SHEEP is used at the recommended dose rate as a single injection, it prevents reinfection with Haemanchus contortus and Teladorsagia (Ostertagia) circumcinata for not less than 91 days, and Tirichstrongylus colubriformis for up to 49 days. WITHHOLDING PERIODS: Macquarie Park, NSW, 2113
Tel: (02) 9889 5400 Product Support Tel: 1800 44 88 92 APVMA: 80572/101142 MEAT: DO NOT USE less than 91 days before slaughter for human consumption. Registered trademark of Nutrien Ag Solutions Limiter

70 mm

**ROLL DIRECTION FIG 5.** 

	PLEASE CHEC	CK I HIS PK	OOF INDROUGE	ILY AS FINAL	. KESPUNS	DIBILLITY FUR	ACCURACT RE	:515 WIIH 100.	
	Check:	1. Content Comments:		☐ Approved, OK to proceed		d Whilst all d	Whilst all care has been taken to present this proof and		
	☐ 1. Content			Name:		••	it's details correctly, it is the customers responsibility to ensure that all details meet their requirements. Your <b>signature indicates your acceptance that all is</b>		
	☐ 2. Colours			Sign:	•••••				
3. Barcode (Numbers)			Date:			<b>correct</b> . If not, please return proof unsigned and with corrections clearly marked.			
	A Pacific	Labels	F/O No <b>34626B</b>	ART VERSION	04 DATE	18.6.2020		Pre Press: D. Harper	
	<b>A.B.N 93 000 881 42</b> 2 Lucca Road, Wyon	<b>25</b> ng NSW 2259	BLACK						
	Ph: (02) 4359 7000 Fax: Email: sales@pacific	. ,	Die-line only (not prin	ted)	PI FΔSI	F NOTE: Colours	s may vary up to 10%	from printed proof	