

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

SWISH

POUR-ON CATTLE LICE AND FLY TREATMENT

ACTIVE CONSTITUENT: 15.0g/L DELTAMETHRIN

For control of Stable Fly (*Stomoxys calcitrans*),
House Fly (*Musca domestica*) and
Lice (*Bovicola bovis*, *Linognathus vituli* and
Haematopinus eurysternus) on beef and dairy cattle.
For control of synthetic pyrethroid susceptible
Buffalo Fly (*Haematobia irritans exigua*)
for up to 3 weeks and for control of
Stable Fly (*Stomoxys calcitrans*)
for up to 1 week on horses.



CATTLE: ZERO (0) WITHHOLDING PERIOD
FOR MILK AND MEAT
ESI (Export Slaughter Interval)
- 21 DAYS



5L

dalgetyah.nutrien.com.au

SWISH

POUR-ON CATTLE LICE AND FLY TREATMENT

DIRECTIONS FOR USE

Restraints
NOT TO BE USED on horses that may be slaughtered for human consumption.

CONTRAINDICATIONS: This product is contraindicated for use on animals that are severely stressed or in ill health. This product should not be used on horses' head forward of the ears. This product should not be applied as a single spot.

PRECAUTION: Facial skin contact may cause temporary tingling/burning.

DOSE RATE FOR CATTLE: 5mL per 100kg liveweight.
The following table may be used as a guide.

Pests Controlled	Rate	Live Body Weight	Critical Comments
Cattle Lice (<i>Bovicola bovis</i>) (<i>Linognathus vituli</i>) (<i>Haematopinus eurysternus</i>)	5 mL	Up to 100kg	Apply to animals in a race or drainage pen as a single strip along the backbone. Retreat later in the season if reinfestation occurs.
	10mL	101-200kg	
	15mL	201-300kg	
	20mL	301-400kg	
Buffalo Fly (<i>Haematobia irritans exigua</i>) Stable Fly (<i>Stomoxys calcitrans</i>) House Fly (<i>Musca domestica</i>)	25mL	401-500kg	Apply to the whole herd – retreatment may be necessary.
	30mL	501-600kg	
	For cattle over 600kg apply 5mL per 100kg liveweight		

DOSE RATE FOR HORSES: 5mL per 100kg liveweight.
The following table may be used as a guide.

Pests Controlled	Rate	Live Body Weight	Critical Comments
Buffalo Fly (<i>Haematobia irritans exigua</i>)	5 mL	Up to 100kg	Using the correct measured dose, apply to horses as two single stripes. Apply approximately 2/3 of measured dose to the midline mane area (from poll to wither) and remaining 1/3 to midline over the rump.
	10mL	101-200kg	
Stable Fly (<i>Stomoxys calcitrans</i>)	15mL	201-300kg	When applying product keep away from head forward of the ears. If product is accidentally splashed into horse's eyes, the eyes should be washed thoroughly as soon as possible, using plenty of water. Contact a veterinary surgeon for advice. Apply to the whole herd – retreatment may be necessary.
	20mL	301-400kg	
House Fly (<i>Musca domestica</i>)	25mL	401-500kg	
	30mL	501-600kg	
For horses over 600kg apply 5mL per 100kg liveweight			

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS:

Cattle: MEAT Zero (0) days; MILK Zero (0) days
Horses: NOT TO BE USED on horses that may be slaughtered for human consumption.

SWISH

POUR-ON CATTLE LICE AND FLY TREATMENT

TRADE ADVICE

Export Slaughter Interval (ESI): DO NOT USE less than 21 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from the distributor on 1300 255 347 or the APVMA website: www.apvma.gov.au/residues/

GENERAL INSTRUCTIONS:

Cattle: To avoid operator insecticide contact when concurrent treatments are carried out, SWISH Pour-On Cattle Lice and Fly Treatment should be applied last. Operators must wear gloves. In a proportion of treated animals, some scurfiness may develop at the site of application. In most cases this scurfiness will disappear within three weeks of application of this product. Apply with recommended applicator. The efficacy of this product was not affected by moderate rainfall when tested against cattle lice.

Horses: Treatment with SWISH will alleviate fly annoyance related behaviour including skin flicking, head and tail twitching and feet stamping for up to 7 days. No adverse reactions were reported in horses treated with deltamethrin at 5 times the recommended dose rate.

It is recommended that the product be allowed to dry thoroughly after application before handling the animal. However, if handling or grooming the animal immediately after application is necessary wear personal protective equipment as indicated in the safety directions on this label.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

DANGEROUS TO FISH. Do not contaminate dams, rivers or streams with the pesticide or used containers.

AVOID CONTACT WITH FOOD.

SAFETY DIRECTIONS

Product is poisonous if swallowed. May irritate the eyes and skin. Facial skin contact may cause temporary facial numbness. Avoid contact with eyes and skin. When using the product wear elbow length PVC gloves. If product on skin, immediately wash area with soap and water. After use and before eating, drinking, or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131 126. Additional information is available in the Material Safety Data Sheet.

Nutrien Ag Solutions Limited
Level 5, Building A, 26 Talavera Road,
Macquarie Park, NSW 2113.
Tel: (02) 9889 5400
Product Support Tel: 1800 44 88 92

DISPOSAL

This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced but may be taken separately.

STORE below 30°C (Room Temperature) in original container, tightly closed in a safe place.

CONDITIONS OF SALE: The use of this product is beyond the control of Nutrien Ag. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Nutrien Ag accepts no responsibility for any consequences resulting from the use of this product. Nutrien Ag will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

IN A TRANSPORT EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE	FOR SPECIALIST ADVICE 1800 033 111 24 HOURS AUSTRALIA WIDE
---	---

APVMA Approval No.: 64199/62005

B:
Exp:



Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: Combustible liquid - Keep away from heat, hot surfaces, sparks, open flames and other ignition sources; No smoking. May damage fertility or the unborn child; May cause respiratory irritation; Do not breathe spray; If inhaled move person to fresh air and keep comfortable for breathing. Store locked up.

230 mm

470 mm

ROLL DIRECTION FIG 5. →

PLEASE CHECK THIS PROOF THOROUGHLY AS FINAL RESPONSIBILITY FOR ACCURACY RESTS WITH YOU.

Check:

- 1. Content
- 2. Colours
- 3. Barcode (Numbers)

Changes & re-submit

Comments: _____
Name: _____
Sign: _____
Date: _____

Approved, OK to proceed

Name: _____
Sign: _____
Date: _____

Whilst all care has been taken to present this proof and it's details correctly, it is the customers responsibility to ensure that all details meet their requirements. Your signature indicates your acceptance that all is correct. If not, please return proof unsigned and with corrections clearly marked.

Pacific Labels
A.B.N 93 000 881 425
2 Lucca Road, Wyong NSW 2259
Ph: (02) 4359 7000 Fax: (02) 4351 1320
Email: sales@pacificlabels.com.au

F/O No 26371

ART VERSION 03

DATE 9.6.2020

Pre Press: D.Harper

BLACK

MAGENTA

CYAN

YELLOW

Die-line only (not printed)

PLEASE NOTE: Colours may vary up to 10% from printed proof.