

READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY



DEFEND™

SPRAY-ON

SHEEP BLOWFLY TREATMENT



250 mm

ACTIVE CONSTITUENT: 50 g/L DICYCLANIL

For the protection of sheep, either off-shears or with any length wool, against strike by dicyclanil-susceptible strains of blowflies (*Lucilia cuprina*) for 18 to 24 weeks.

For the protection of mulesing and marking wounds on sheep against strike by dicyclanil-susceptible strains of blowflies (*Lucilia cuprina*) during the wound healing process.

DEFEND Spray-On contains dicyclanil, a member of the pyrimidine class of chemicals.

It is effective against sensitive strains of blowfly (*Lucilia cuprina*).

Resistance may develop to any chemical.



Dalgety
Animal Health

20 L

dalgetyah.nutrien.com.au

141255_0119

200 mm

← ROLL DIRECTION FIG 6.

PLEASE CHECK THIS PROOF THOROUGHLY AS FINAL RESPONSIBILITY FOR ACCURACY RESTS WITH YOU.

Check:

- 1. Content
- 2. Colours
- 3. Barcode (Numbers)

Changes & re-submit

Comments:.....

.....

.....

.....

Approved, OK to proceed

Name:.....

Sign:

Date:.....

Whilst all care has been taken to present this proof and it's details correctly, it the customers responsibility to ensure that all details meet their requirements. Your signature indicates your acceptance that all is correct. If not, please return proof unsigned and with corrections clearly marked.

 **Pacific Labels**

A.B.N 93 000 881 425
2 Lucca Road, Wyong NSW 2259
Ph: (02) 4359 7000 Fax: (02) 4351 1320
Email: sales@pacificlabels.com.au

F/O No 39036A

ART VERSION 02

DATE 12.3.2020

Pre Press: D. Harper

BLACK

MAGENTA

PMS 021

PMS 292

CYAN

YELLOW

PMS 364

Die-line only (not printed)

PLEASE NOTE: Colours may vary up to 10% from printed proof.

DEFEND™ SHEEP BLOWFLY SPRAY-ON TREATMENT




DIRECTIONS FOR USE

Restraints: DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

DOSAGE AND ADMINISTRATION – SHAKE WELL BEFORE USE

DEFEND Spray-On is applied as a low volume treatment on to areas where protection is required. When applying DEFEND, the nozzle of the applicator gun should be high enough above the fleece (approximately 15-25 cm), to achieve a 15 cm wide band at each pass for protection of body strike. For treatment of mulesing wounds, the band must be wide enough to cover the wound and 25 mm onto the surrounding wool. DO NOT apply during gusty or windy weather as spray drift may reduce the period of protection. It is good agricultural practice for mulesing/ marking to be carried out at times of minimal fly activity. Reduced efficacy may occur in sheep affected with dermatophilosis.

FOR THE PREVENTION OF BODY STRIKE (Diagram 1)					CRITICAL COMMENTS
SHEEP WEIGHT (kg)	TOTAL VOLUME / SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS	NO. OF SHEEP TREATED PER 20L	
10-20	14	7	2	1428	<ol style="list-style-type: none"> Use only a spray-on applicator gun recommended by Landmark. Ensure the applicator gun is clean. Calibrate applicator gun using an accurate volume measure. Ensure all areas to be protected are treated. When preventing body strike ensure bands overlap along the midline – see Diagram 1. When protecting mulesing wound from fly strike, take special care to treat the wound and the surrounding 25 mm of wool – see Diagram 2. When protecting marking wounds from fly strike, take special care to treat the wound and surrounding wool – see Diagram 3. This product is not recommended for treating existing strikes. Use a product registered as a dressing for existing strikes. Heavy rain following application of this product could diminish the period of protection. DEFEND Spray-on contains a scourable colouring agent for easy band identification. Withholding periods must be observed following application of DEFEND for any purpose including after mulesing and/or marking as finite residues will occur.
21-30	16	8	2	1250	
31-50	20	10	2	1000	
>50	24	12	2	833	
FOR THE PREVENTION OF BREACH STRIKE (Diagram 1)					
SHEEP WEIGHT (kg)	TOTAL VOLUME / SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS	NO. OF SHEEP TREATED PER 20L	
10-20	7	7	1	2857	
21-30	8	8	1	2500	
31-50	10	10	1	2000	
>50	12	12	1	1666	
FOR THE PREVENTION OF POLL STRIKE (Diagram 1)					
Rams and sheep susceptible to poll strike should be treated. Apply 5mL of DEFEND Spray-on to the areas requiring protection, around the poll and horns.					
FOR THE PREVENTION OF FLY STRIKE AFTER MULESING (Diagram 2)					
Apply two bands, one over each side of the mulesing wound and overlapping 25mm onto the surrounding wool.					
SHEEP WEIGHT (kg)	TOTAL VOLUME / SHEEP (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS	NO. OF SHEEP TREATED PER 20L	
Up to 30	8	4	2	2500	
31-50	10	5	2	2000	
FOR THE PREVENTION OF FLY STRIKE AFTER MARKING (Diagram 3)					
<ul style="list-style-type: none"> When a tail has been amputated, apply 4mL over the crutch ensuring sufficient coverage of the open wound. When the tail has not been amputated, ensure sufficient coverage of the tail where the rubber ring has been applied. In the case of ram lambs a further 4 mL can be sprayed over and around the scrotum area after the rubber ring has been applied.					
LAMBS WEIGHT (< 30 kg)	TOTAL VOLUME / LAMB (mL)	VOLUME PER TREATMENT BAND (mL)	NO. OF BANDS	NO. OF LAMBS TREATED PER 20L	
Ram lambs	8	4	2	2500	
Ewe lambs	4	4	1	5000	



TRADE ADVICE – EXPORT SLAUGHTER INTERVALS (ESIs): SHEEP (long wool) DO NOT use less than 60 days before slaughter for export. SHEEP (6 weeks wool or less) DO NOT USE less than 120 days before slaughter for export. Before using this product, confirm the current ESI from Landmark on 1300 255 347 or the APVMA website (www.apvma.gov.au/residues). Use of this product may result in wool residues that may not comply with European Union environmental quality standards. Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.

SHAKE WELL BEFORE USE

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT: DO NOT contaminate dams, rivers, or streams with the product or used containers.

SAFETY DIRECTIONS: May irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and using the product wear elbow length rubber gloves. Wash hands after use.

FIRST AID: If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131 126.

ADDITIONAL USER SAFETY INFORMATION: If additional information is required, refer to the Safety Data Sheet. For a copy phone 1300 255 347.

ENVIRONMENTAL PROTECTION: DO NOT contaminate dams, rivers, streams or other waterways with the chemical or used container.

DISPOSAL: Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. DO NOT dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

STORAGE: KEEP OUT OF REACH OF CHILDREN. Store below 30°C (Room Temperature). Protect from frost. Store in original container tightly closed in a dry, cool place. DO NOT store in direct sunlight. DO NOT re-use the container.

CONDITIONS OF SALE: The use of this product is beyond the control of Nutrien Ag Solutions. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Nutrien Ag Solutions accepts no responsibility for any consequences resulting from the use of this product. Nutrien Ag Solutions will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: The product is not classified according to the Globally Harmonised System (GHS).

Nutrien Ag Solutions

Suite 3, Level 1, Building B, 11 Talavera Road, Macquarie Park, NSW, 2113 • Tel: (02) 9889 5400

Product Support Tel: 1800 44 88 92

APVMA Approval No.: 65651/117850 141256_0119



B:
Expiry:

9 331984 679571

IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE

FOR SPECIALIST ADVICE IN AN EMERGENCY DIAL
1800 033 111
24 HOURS AUSTRALIA WIDE

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

GENERAL DIRECTIONS

DEFEND Spray-On provides season long protection to sheep against body, breach and poll strike and may be used either off-shears or on sheep with any length wool. DEFEND Spray-On also protects mulesing and marking wounds on lambs against fly strike during the wound healing process. The dose volume is determined by the weight of sheep being treated. Apply DEFEND Spray-On according to the diagrams and tables in the Directions for Use. DO NOT dilute. DO NOT mix with any other product. Apply directly from the container or an appropriate back pack using a spray-on applicator gun recommended by Landmark. DEFEND Spray-On may be applied to all breeds of sheep to obtain protection against fly strike.

Significant soiling with dags, mud, dust or urine may reduce efficacy. Where significant dags are present, a light crutching is indicated prior to treatment. Protection period may be reduced if sheep scour heavily following application.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 28 days before slaughter for human consumption.

MILK: DO NOT USE on ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

WOOL HARVEST INTERVAL: DO NOT USE less than 3 months before shearing or fibre collection.

200 mm

ROLL DIRECTION FIG 5. →

PLEASE CHECK THIS PROOF THOROUGHLY AS FINAL RESPONSIBILITY FOR ACCURACY RESTS WITH YOU.

Check:

1. Content Changes & re-submit Approved, OK to proceed

2. Colours Comments:..... Name:.....

3. Barcode (Numbers) Sign:

..... Date:

Whilst all care has been taken to present this proof and it's details correctly, it the customers responsibility to ensure that all details meet their requirements. Your signature indicates your acceptance that all is correct. If not, please return proof unsigned and with corrections clearly marked.

Pacific Labels

A.B.N 93 000 881 425
2 Lucca Road, Wyong NSW 2259
Ph: (02) 4359 7000 Fax: (02) 4351 1320
Email: sales@pacificlabels.com.au

F/O No 39036B

ART VERSION 02

DATE 12.3.2020

Pre Press: D. Harper

BLACK

PMS 364

PMS 485

CYAN

YELLOW

MAGENTA

Die-line only (not printed)

PLEASE NOTE: Colours may vary up to 10% from printed proof.

250 mm