


CAUTION
 KEEP OUT OF REACH OF CHILDREN
 FOR ANIMAL TREATMENT ONLY
 READ SAFETY DIRECTIONS BEFORE OPENING OR USING

PROXIMAX POUR-ON FOR BEEF AND DAIRY CATTLE



ACTIVE CONSTITUENT:
 5 mg/ml Eprinomectin

For the treatment and control of internal and external parasites of beef and dairy cattle and internal parasites of deer.

Nutrien Ag Solutions Limited
 Suite 3, Level 1, Building B
 11 Talavera Road, Macquarie Park, NSW, 2113
 Tel: (02) 9889 5400
 Product Support Tel: 1800 44 88 92



Dalgety
 Animal Health

20L

dalgetyah.nutrien.com.au

105587_0420

180 mm

ROLL DIRECTION FIG 5.

180 mm

PLEASE CHECK THIS PROOF THOROUGHLY AS FINAL RESPONSIBILITY FOR ACCURACY RESTS WITH YOU.

Check:

- 1. Content
- 2. Colours
- 3. Barcode (Numbers)

Changes & re-submit

Comments:.....

Approved, OK to proceed

Name:.....
 Sign:
 Date:.....

Whilst all care has been taken to present this proof and it's details correctly, it is the customers responsibility to ensure that all details meet their requirements. Your signature indicates your acceptance that all is correct. If not, please return proof unsigned and with corrections clearly marked.

Pacific Labels

A.B.N 93 000 881 425
 2 Lucca Road, Wyong NSW 2259
 Ph: (02) 4359 7000 Fax: (02) 4351 1320
 Email: sales@pacificlabels.com.au

F/O No 40791F ART VERSION 02 DATE 7.5.20

Pre Press: G.A.Davis

BLACK	MAGENTA		PMS 021
CYAN	YELLOW		

Die-line only (not printed)

PLEASE NOTE: Colours may vary up to 10% from printed proof.

180 mm

PROXIMAX POUR-ON FOR BEEF AND DAIRY CATTLE

INDICATIONS:

For the treatment and control of gastrointestinal roundworms (including inhibited immature *Ostertagia ostertagi*) and lungworm in cattle.

For sucking and biting lice, sarcoptic and chorioptic mange mites and buffalo fly in beef and dairy cattle.

Proximax Pour-On aids in control of cattle tick (*Boophilus microplus*). For the treatment and control of gastrointestinal roundworms and lungworm in deer.

PROXIMAX Pour-On for Beef and Dairy Cattle contains eprinomectin, a member of the macrocyclic lactone family of drenches. It is effective against eprinomectin sensitive strains of internal parasites. Resistance can develop to any drench.

CATTLE: Gastrointestinal Roundworms:

For the control of adult and immature:

Ostertagia ostertagi (including inhibited L4) – small brown stomach worm, *O. lyrata* – brown stomach worm (adults only)
Ostertagia spp. *Haemonchus placei* – barbers pole worm, *Trichostrongylus axei* – stomach hair worm, *T. colubriformis* – intestinal hair worm, *Trichostrongylus spp.* *Cooperia spp.* (including inhibited L4) – small intestinal worm, *C. oncophora*, *C. pectinata*, *C. punctata*, *C. sumabada*, *Oesophagostomum radiatum* – nodule worm, *Oesophagostomum spp.*

Nematodirus helvetianus – thin necked intestinal worm
Bunostomum phlebotomum – hookworm, *Trichuris spp.* – whip worm (adults only)

Lungworm Adult and immature *Dictyoacaulus viviparus*

Sucking and biting lice *Linognathus vituli*, *Haematopinus eurysternus*

Bovicola (Damalinea) bovis, *Solenopotes capillatus*

Mites *Choripetes bovis*, *Sarcoptes scabiei var bovis*

Buffalo Fly *Haematobia irritans exigua*

When used as directed, Proximax Pour-On also aids control of:

Cattle tick *Rhipicephalus (Boophilus) microplus*

(not recommended for use in strategic tick control programs)

PERSISTENT ACTIVITY IN CATTLE:

Proximax Pour-On when given at the recommended dosage of 1 mL/10 kg liveweight controls infections with:

Parasite*	Prolonged Activity
<i>Ostertagia spp.</i>	Up to 28 days
<i>Cooperia spp.</i>	Up to 28 days
<i>Dictyoacaulus viviparus</i>	28 days
<i>Nematodirus helvetianus</i>	Up to 28 days
<i>Oesop hagostomum radiatum</i>	Up to 28 days
<i>Haemonchus placei</i>	21 days
<i>Trichostrongylus spp.</i>	21 days

*The following parasite species are included within each of the relevant genera: *Ostertagia ostertagi*, *O. leptospicularis*, *O. lyrata*, *Cooperia oncophora*, *C. punctata*, *C. sumabada*, *Trichostrongylus axei*, *T. colubriformis*.

DEER – Gastrointestinal roundworms:

Ostertagia – type spp. *(adults)

Trichostrongylus spp. (adults and fourth stage larvae)

Mazamastrongylus spp. (adults)

Oesophagostomum spp. (adults)

* includes *O. mossi/dikmansi*, *O. lyrata*, *O. leptospicularis*, *Spiculoptera spiculoptera*, *Sp. asymmetrica* and *Aptergia quadrispiculata*

Lungworms:

Dictyoacaulus viviparus (adults and fourth stage larvae). For best results Proximax Pour-On for Beef and Dairy Cattle should be part of a total parasite control program including internal and external parasites. Resistance may develop to any drench Consult your veterinarian for a complete parasite control program.

DIRECTIONS FOR USE:

DOSAGE AND ADMINISTRATION:

Proximax Pour-On for Beef and Dairy Cattle contains eprinomectin, a member of the macrocyclic lactone family of drenches. It is effective against eprinomectin sensitive strains of internal parasites. Resistance can develop to any drench.

For best results PROXIMAX Pour-On should be part of a total parasite control program including internal and external parasites. Resistance can develop to any drench.

Consult your veterinarian for a complete parasite control program.

A representative sample of animals should be weighed with either scales or a weighband. Dose the mob accordingly to the heaviest animal by liveweight in the group (cows, bulls, calves, heifers, hinds, stags, fawns etc). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

This product is formulated only for pour-on application to cattle and deer; do not use in other species. The formulation should be applied along the topline in a narrow strip extending from the withers to the tail head. The dose rate is 1 mL for each 10 kg liveweight.

A 20L drum contains enough solution to treat 800 head of 250kg cattle or 2000 head of 100kg deer.

Liveweight (kg)	Dose Volume (mL)	Liveweight (kg)	Dose Volume (mL)
Up to 50	5	251 - 300	30
51-75	7.5	301 - 350	35
76-100	10	351 - 400	40
101-125	12.5	401 - 450	45
126-150	15	451 - 500	50
151 - 175	17.5	501 - 550	55
176 - 200	20	551 - 600	60
201 - 250	25	601 - 650	65

Heavier animals (for example mature bulls) should receive an additional 5 mL for each 50 kg over 650 kg.

WITHHOLDING PERIODS:

CATTLE MILK: Zero (0) days

CATTLE MEAT/DEER MEAT: Zero (0) days

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL

EXPORT SLAUGHTER INTERVAL (ESI): Zero (0) days.

Before using this product confirm the current ESI from Landmark Operations Limited or the APVMA website (www.apvma.gov.au/residues). Consult your veterinarian for the assistance in the diagnosis, treatment and control of parasitism.

ANIMAL SAFETY

Studies have demonstrated a wide safety margin. Five times the recommended use level had no adverse effect on cattle or deer. Three times the recommended use level had no adverse effect on breeding performance of cows and bulls.

NOTE TO USER

Rainfall before or after treatment will not affect the efficacy of Proximax Pour-On.

PRECAUTIONS

Close container when not in use.

Do not apply to areas of the backline covered with mud or manure.

This product is for application to skin surface only, do not give orally or by injection.

INSTRUCTIONS FOR THE 20L PACK:

- Replace the shipping cap with the dip tube assembly.
- Attach the draw-off tubing to the dosing applicator and to the top stem on the dip tube cap.
- Gently prime the dosing applicator checking for leaks.
- Follow the dosing applicator manufacturer's directions for adjusting the dose and proper use and maintenance of the equipment.

STORAGE

Store below 30°C (Room Temperature).

Store bottle or pack in carton to protect from light.

DISPOSAL

If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar management program site. The cap should not be replaced, but may be recycled separately with the container. Triple pressure rinse Proximax Pour-On containers and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken and crushed or punctured containers 500 mm below the surface and in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and trees roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

ENVIRONMENT PROTECTION

Studies indicate that when eprinomectin comes in contact with the soil, it readily and tightly binds to the soil and becomes inactive over time. Containers and any residual contents should be disposed of safely in an approved landfill as free eprinomectin may adversely affect fish and certain water-borne organisms.

SAFETY DIRECTIONS

Poisonous if swallowed. Will irritate the eyes. Avoid contact with eyes. Repeated exposure may cause allergic disorders. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. For further information, refer to the Safety Data Sheet which is available from the supplier.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which can be obtained from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of Nutrien Ag Solutions Limited. Any provisions or rights under the Australian Consumer Law which cannot be excluded by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies expressed or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application of this product are excluded. Nutrien Ag Solutions Limited does not accept any liability (including consequential loss and/or negligence) for any loss or damage connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.

APVMA Approval No: 87053/117447

Nutrien Ag Solutions Limited

Suite 3, Level 1, Building B,
11 Talavera Road, Macquarie Park, NSW, 2113,
Tel: (02) 9889 5400 Product Support Tel: 1800 44 88 92

Not classified as Hazardous according to the Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia criteria.



BATCH
EXPIRY

IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE

FOR SPECIALIST ADVICE IN AN EMERGENCY DIAL
1800 033 111
24 HOURS AUSTRALIA WIDE

105587_0420

ROLL DIRECTION FIG 5. →

180 mm

PLEASE CHECK THIS PROOF THOROUGHLY AS FINAL RESPONSIBILITY FOR ACCURACY RESTS WITH YOU.

Check:

1. Content

Changes & re-submit

Comments:.....

2. Colours

.....

3. Barcode (Numbers)

.....

Approved, OK to proceed

Name:.....

Sign:

Date:

Whilst all care has been taken to present this proof and it's details correctly, it is the customers responsibility to ensure that all details meet their requirements. Your signature indicates your acceptance that all is correct. If not, please return proof unsigned and with corrections clearly marked.

Pacific Labels

A.B.N 93 000 881 425
2 Lucca Road, Wyong NSW 2259
Ph: (02) 4359 7000 Fax: (02) 4351 1320
Email: sales@pacificlabels.com.au

F/O No 40791B ART VERSION 02 DATE 7.5.20

BLACK

PMS 021

Die-line only (not printed)

PLEASE NOTE: Colours may vary up to 10% from printed proof.

Pre Press: G.A.Davis