

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
FOR ANIMAL TREATMENT ONLY

# CENTURIION® POUR-ON LICE TREATMENT FOR SHEEP



ACTIVE CONSTITUENT: 35 g/L IMIDACLOPRID  
SOLVENT: 300 g/L N-METHYL-2-PYRROLIDONE  
For the control of neonicotinoid susceptible body lice  
*Bovicola ovis* on shorn sheep up to 7 days off-shears and  
unshorn lambs up to 2 months of age.  
This product can be expected to protect sheep  
against re-infestation with lice for 4 weeks after  
treatment when applied within 24 hrs of shearing.



**Dalgety**  
Animal Health

20 L

[dalgetyah.nutrien.com.au](http://dalgetyah.nutrien.com.au)

105478\_0320

180 mm

180 mm

ROLL DIRECTION FIG 5. →

**PLEASE CHECK THIS PROOF THOROUGHLY AS FINAL RESPONSIBILITY FOR ACCURACY RESTS WITH YOU.**

**Check:**

- 1. Content
- 2. Colours
- 3. Barcode (Numbers)

Changes & re-submit

Comments: \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

Approved, OK to proceed

Name: \_\_\_\_\_  
Sign: \_\_\_\_\_  
Date: \_\_\_\_\_

Whilst all care has been taken to present this proof and it's details correctly, it is the customers responsibility to ensure that all details meet their requirements. Your **signature indicates your acceptance that all is correct**. If not, please return proof unsigned and with corrections clearly marked.



A.B.N 93 000 881 425  
2 Lucca Road, Wyong NSW 2259  
Ph: (02) 4359 7000 Fax: (02) 4351 1320  
Email: [sales@pacificlabels.com.au](mailto:sales@pacificlabels.com.au)

F/O No 38594F ART VERSION 01 DATE 8.4.2020

Pre Press: D. Harper

BLACK	MAGENTA	PMS 021	PMS 124
CYAN	YELLOW	PMS 348	

Die-line only (not printed)

PLEASE NOTE: Colours may vary up to 10% from printed proof.

**DIRECTIONS FOR USE**

This product can be used on all adult sheep up to 7 days off-shears and on unshorn lambs up to 2 months of age. Apply the entire dose to sheep under 30 kg as a single broadband stripe from poll to tail butt. For sheep over 30 kg, apply half the dose as a broadband stripe either side of the midline from the mid neck to the butt of the tail. The stripes should be no further than 10cm apart. Refer to the diagrams below. A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by liveweight in each group (ewes, wethers, rams, lambs). DO NOT underdose. Where there is a large variation in size within the group (e.g. lambs), dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

Bodyweight (kg)	Dose (mL)	Method	Apply in the pattern below
Below 6kg	8	Single Stripe	
6.0-8.0	12		
8.1-10	15		
10.1-12.5	20		
12.6-15	25		
15.1-20	30		
20.1-30	40	Double Stripe	
30.1-55	60		
55.1-80	80		
80.1-100	90		

Add 10mL for each 10kg above 100kg.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**GENERAL INSTRUCTIONS**

Apply product only with a wide band or multi-stripe applicator. To prevent reinfestation treated sheep must not be mixed with lice free sheep until 6 weeks after treatment. Live lice on ewes treated within 6 weeks of lambing can infect lambs from these ewes. Sheep must be treated within 7 days of shearing. Unshorn lambs can be treated up to 2 months of age. Efficacy is not affected by moderate rainfall either before or after treatment (50 mm rain 1 hour prior to treatment and/or 12.5 mm 2 hours after treatment and/or 25 mm 24 hours after treatment). Treated sheep must not be mixed with lousy sheep. DO NOT use on sheep which have not been cleanly shorn for any reason e.g., mycotic dermatitis. Management practices which promote good lice control such as complete musters, accurate treatment of every animal according to label, ensuring all animals can be treated at the same time, no split shearings, stock-proof fencing to prevent external introductions, quarantine and lice treatment of all newly purchased stock are essential for effective lice control. Treated sheep could potentially develop backline dermatitis following prolonged exposure to extreme heat and sunlight without access to shade. DO NOT leave container, hose or applicator in the sun for long periods of time. Centurion® Pour-On Lice Treatment For Sheep may be used in conjunction with worm control drenches and at the same time as vaccination. Centurion® Pour-On Lice Treatment For Sheep contains a blue dye to identify treated sheep. This dye is scourable from the wool. The product can be detrimental to certain plastics, paints and surface finishes. Avoid contact with such surfaces. Centurion® Pour-On Lice Treatment For Sheep is safe for use on breeding rams.

**WITHHOLDING PERIODS**

**MEAT: DO NOT USE less than 21 days before slaughter for human consumption.**

**MILK: DO NOT USE on female sheep which are producing, or may in the future produce milk or milk products for human consumption.**

**WOOL: DO NOT USE less than 2 months before shearing or fibre collection.**

**EXPORT SLAUGHTER INTERVAL (ESI):** DO NOT slaughter for export less than 63 days after treatment. The ESI on this label was correct at the time of label approval. Before using this product confirm the current ESI from either Nutrien Ag Solutions Limited (1800 448 892) or the APVMA website [www.apvma.gov.au/residues](http://www.apvma.gov.au/residues).

**SAFETY DIRECTIONS:** Will irritate the eyes. Avoid contact with eyes. When using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and washable hat and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands arms and face thoroughly with soap and water. After each day's use, wash goggles and contaminated clothing. **Rehandling Interval:** DO NOT come into contact with treated sheep until the spray has dried unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing).

**FIRST AID:** If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126.

**ENVIRONMENTAL STATEMENTS:** Imidacloprid is highly toxic aquatic species. DO NOT contaminate dams, ponds, rivers, waterways or drains with pesticide or used containers.

**DISPOSAL OF CONTAINERS:** Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. If the container has the *drumMUSTER* logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any *drumMUSTER* collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.


**STORAGE:** Store below 30°C (room temperature) in closed, original container and protect from light.

**SAFETY DATA SHEET:** For further information, refer to the Safety Data Sheet which is available from the supplier.

**CONDITIONS OF SALE:** The use of this product is beyond the control of Nutrien Ag Solutions Limited. Any provisions or rights under the Australian Consumer Law which cannot be excluded by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies expressed or implied under common law, statute or otherwise, in relation to the sale, supply, storage, use or application of this product are excluded. Nutrien Ag Solutions Limited does not accept any liability (including consequential loss and/or negligence) for any loss or damage connected with the sale, supply, storage, use or application of this product except for liability which cannot be excluded by statute.

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: Toxic to aquatic life with long lasting effects. Avoid release to the environment. Collect spillage.

© Registered trademark of Nutrien Ag Solutions



**UN No. 3082**  
**ENVIRONMENTALLY HAZARDOUS**  
**SUBSTANCE, LIQUID, N.O.S.**  
**(Contains Imidacloprid)**

**Nutrien Ag Solutions Limited**

Level 5, Building A, 26 Talavera Road, Macquarie Park, NSW, 2113  
Tel: (02) 9889 5400

APVMA Approval No.: 83012/117552

105585\_0320



Batch Number:

Expiry:

9 331984 703252

**IN A TRANSPORT EMERGENCY**  
**DIAL 000**  
**POLICE OR FIRE BRIGADE**

**FOR SPECIALIST ADVICE IN AN**  
**EMERGENCY DIAL**  
**1800 033 111**  
**24 HOURS AUSTRALIA WIDE**

180 mm

180 mm

← **ROLL DIRECTION FIG 6.**



A.B.N 93 000 881 425

2 Lucca Road, Wyong NSW 2259

Ph: (02) 4359 7000 Fax: (02) 4351 1320

sales@pacificlabels.com.au / www.pacificlabels.com.au

Job No: **38594B** Version: **4** Date: **09.03.2022**

Operator: **Dalys** Client: **Nutrien Ag**

Whilst all care has been taken to avoid mistakes, **THIS IS YOUR PROOF - PROOF READING IS YOUR RESPONSIBILITY.** Please read very carefully as you are 100% responsible for the accuracy once approval has been given. Delays in approval WILL effect the delivery date. Delivery date confirmation can only be advised once your artwork approval has been received

Approved by: \_\_\_\_\_  
Date: \_\_\_\_/\_\_\_\_/\_\_\_\_

**PRINT COLOURS**



**FINISHING**

Material:	PP White	Qty per Roll:	250
Core size:	76 mm		

PLEASE NOTE that PDF files sent via email are not COLOUR CORRECT due to non calibration on your monitor. Colours may vary up to 10% from printed proof. The spot colour swatches shown on this proof are indicative only - all Pantone PMS colours are to be matched against Pantone Scale guides or supplied samples. Colours used for all die cuts, varnish, foil and other embellishments are for position only and not representative of colour or final result.