

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

MOXIMAX[®] LONG ACTING INJECTION FOR CATTLE

ACTIVE CONSTITUENT: 100 mg/mL MOXIDECTIN



For the treatment and control
of moxidectin sensitive internal and
external parasites in cattle.



Dalgety
Animal Health

1P4375F-1

500mL

dalgetyah.nutrien.com.au

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY**

MOXIMAX[®] LONG ACTING INJECTION FOR CATTLE

ACTIVE CONSTITUENT: 100 mg/mL MOXIDECTIN

**BEFORE USING THIS PRODUCT. READ
ALL DIRECTIONS ON THE CARTON.**

DIRECTIONS FOR USE:

DOSAGE AND ADMINISTRATION

Discard unused portion after 6 months of first broaching container.

Dose Rate:

1 mg moxidectin/kg liveweight
(1 mL/100 kg liveweight).

**Retreatment Interval: Do not retreat
with MOXIMAX LONG ACTING
INJECTION FOR CATTLE for at least 56
days after administration.**

WITHHOLDING PERIODS

**MEAT: DO NOT USE less than 56
days before slaughter for human
consumption.**

**MILK: DO NOT USE on lactating cows
or within 80 days of calving where
milk or milk products may be used for
human consumption.**

TRADE ADVICE

**EXPORT SLAUGHTER INTERVAL (ESI):
DO NOT USE less than 108 days before
slaughter for export. The ESI on this
label was correct at the time of label
approval.**

**Before using this product confirm the
current ESI from either Nutrien Ag
Solutions Limited on 1800 44 88 92 or
the APVMA website [www.apvma.gov.
au/residues](http://www.apvma.gov.au/residues)**

**DO NOT USE in cattle that may be
exported live.**

1P4375B-1

FIRST AID:

If poisoning occurs, contact a doctor
or Poisons Information Centre. Phone
Australia 13 11 26. Additional information
is in the Safety Data Sheet (SDS).

DISPOSAL

Dispose of container by wrapping with
paper and putting in garbage.
Discarded needles should immediately be
placed in a designated and appropriately
labelled "sharps" container. The container
should be of a type to reduce the
possibility of injury to handlers during
collection and disposal. Incineration is the
preferred method of disposal, otherwise
sharps should be buried at a suitable site,
such as an on-farm chemical disposal pit
located away from water courses.

STORAGE

Store below 30°C (Room temperature).
Protect from light.
APVMA: 91130/131005

[B]:

EXP:

Nutrien Ag Solutions Limited
Level 5, Building A, 26 Talavera Road,
Macquarie Park, NSW 2113.

Tel: (02) 9889 5400

Support Tel: 1800 44 88 92