

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



**Dalgety**  
Animal Health

# CATTLEMAX<sup>®</sup> PLUS

INJECTION FOR CATTLE

**ACTIVE CONSTITUENTS: 10 g/L IVERMECTIN,  
100 g/L CLORSULON**

For the treatment and control of ivermectin and clorsulon sensitive strains of internal and external parasites of beef and dairy cattle, including adult liver fluke.

Resistance may develop to any chemical.

**INTERNAL AND  
EXTERNAL PARASITES**



[dalgetyah.com.au](http://dalgetyah.com.au)

**500 mL**

**BEFORE USING THIS PRODUCT, READ ALL DIRECTIONS ON THE CARTON.**

**DIRECTIONS FOR USE:**

The recommended dose is 1 mL CATTLEMAX® PLUS INJECTION FOR CATTLE per 50 kg liveweight (equivalent to 200 µg ivermectin and 2 mg clorsulon per kg liveweight) by subcutaneous injection. This product is not to be used intravenously or intramuscularly.

**WITHHOLDING PERIODS:**

**MEAT: DO NOT USE LESS than 28 days before slaughter for human consumption.**  
**MILK: NIL**

**TRADE ADVICE: Export Slaughter Interval (ESI): DO NOT USE less than 42 days before slaughter for export.** The ESI on this label was correct at the time of label approval. Before using this product confirm the current ESI from either

Loveland Agri Products Pty Ltd on 1800 448 892 or the APVMA website [www.apvma.gov.au/residues.shtml](http://www.apvma.gov.au/residues.shtml)

**SAFETY TO CATTLE:**

Studies have demonstrated a wide margin of safety. Pregnant and breeding cattle may be treated provided normal care is taken in handling.

**SAFETY DIRECTIONS:**

Poisonous if swallowed. Avoid contact with eyes, skin and clothing.

**FIRST AID:** If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

**ADDITIONAL USER SAFETY:**

Additional information is listed in the Safety Data Sheet, available by request.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT:**

Ivermectin is extremely toxic to aquatic species. Do not contaminate dams,

rivers, streams or other waterways with the chemical or used container.

**DISPOSAL:** Dispose of empty container and packaging by wrapping with paper and putting in garbage. Discarded needles/sharps should immediately be placed in a designated and appropriately labelled 'sharps' container.

**STORAGE:** Store below 30 °C (room temperature). STORE BOTTLE IN CARTON TO PROTECT FROM LIGHT.

**Loveland Agri Products Pty Ltd**

Level 5, Building A,  
26 Talavera Road,  
Macquarie Park,  
NSW, 2113

Tel: (02) 9889 5400  
Product Support  
Tel: 1800 44 88 92  
[dalgetyah.com.au](http://dalgetyah.com.au)

APVMA Approval No.:  
90406/128773

Batch Number:

Expiry:

**SAFETY TO CATTLE:** Studies have demonstrated a wide margin of safety. Pregnant and breeding cattle may be treated provided normal care is taken in handling.

**Caution: avoid carcass damage**

1. Sterilise all injection apparatus by boiling (or equivalent) before use. Avoid use of strong disinfectants on apparatus.
2. Maintain cleanliness at all times.
3. Keep needles sharp and clean. Replace frequently.
4. Use shortest needle possible, certainly not exceeding 15 mm.
5. As far as possible avoid injection of animals during wet weather or under dusty conditions.
6. This product should be injected only under the skin.
7. If possible inject high on the neck behind the ear.

**SAFETY DIRECTIONS:** Poisonous if swallowed. Avoid contact with eyes, skin and clothing.

**FIRST AID:** If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

**ADDITIONAL USER SAFETY:** Additional information is listed in the Safety Data Sheet, available by request.

**ENVIRONMENTAL PROTECTION:** Ivermectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other watercourses with the chemical or used container.

**DISPOSAL:** Dispose of empty container and packaging by wrapping with paper and putting in garbage. Discarded needles/sharps should immediately be placed in a designated and appropriately labelled 'sharps' container.

**STORAGE:** Store below 30 °C (room temperature). STORE BOTTLE IN CARTON TO PROTECT FROM LIGHT.

**CONDITIONS OF SALE:** This product should only be used in accordance with the label and manufacturer's instructions. Product performance may be affected by factors which are beyond the control of Loveland Agri Products Pty Ltd and its

Associates (as that term is defined in the Corporations Act 2001 Cth) (Associates). If symptoms persist, please consult your veterinarian or animal health specialist. In no case shall Loveland Agri Products Pty Ltd or its Associates be liable for consequential, incidental, special or exemplary damages or losses.

All sales (wholesale and retail) are governed by the General Terms and Conditions of Sale available at [lovelandagriproducts.com.au](http://lovelandagriproducts.com.au) or from your retailer upon request. These terms can only be varied by Loveland Agri Products Pty Ltd.

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: Suspected of causing cancer. May damage fertility or the unborn child. Harmful if swallowed. Harmful to aquatic life with long lasting effects. Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Wash thoroughly after handling. Do not eat, drink or smoke when using this product. Avoid release to the environment. Wear protective gloves/ protective clothing/eye protection/face protection/hearing protection. IF SWALLOWED: Call a POISON CENTER/doctor if you feel unwell. Rinse mouth. IF exposed or concerned: Get medical advice/attention. Store locked up. Dispose of contents/container in accordance with local/regional/national regulations. Product contains: Ivermectin (10 g/L) and Clorsulon (100 g/L)

IN A TRANSPORT EMERGENCY  
**DIAL 000**  
POLICE OR FIRE BRIGADE

FOR SPECIALIST ADVICE  
IN AN EMERGENCY  
**1800 033 111**  
24 HOURS AUSTRALIA WIDE

**Loveland Agri Products Pty Ltd**

Level 5, Building A, 26 Talavera Road, Macquarie Park, NSW, 2113

Tel: (02) 9889 5400 • Product Support Tel: 1800 44 88 92

[dalgetyah.com.au](http://dalgetyah.com.au)

APVMA Approval No.: 90406/128773

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**CATTLEMAX<sup>®</sup>**  
**PLUS**

INJECTION FOR CATTLE

**ACTIVE CONSTITUENTS: 10 g/L IVERMECTIN, 100 g/L CLORSULON**

For the treatment and control of ivermectin and clorsulon sensitive strains of internal and external parasites of beef and dairy cattle, including adult liver fluke.

CATTLEMAX® PLUS INJECTION FOR CATTLE contains ivermectin, a member of the macrocyclic lactone family of chemicals and clorsulon, a member of the sulphonamide family of chemicals. When used at the recommended dose level of 200 µg ivermectin/kg and 2 mg clorsulon/kg liveweight, it is effective against sensitive strains of the following internal parasites:

Resistance may develop to any chemical.

#### GASTRO-INTESTINAL ROUNDWORMS:

##### Adult and immature:

*Ostertagia ostertagi* - Small brown stomach worms (including third and fourth stage larvae and inhibited immatures)

*Haemonchus placei* - Barber's pole worm

*Trichostrongylus axei* - Stomach hair worm (including third and fourth stage larvae)

*Trichostrongylus colubriformis* - Intestinal hair worm

*O. lyrata* - Stomach worm

*Cooperia* spp - Small intestinal worm (including third and fourth stage larvae)

*Oesophagostomum radiatum* - Nodule worm (including third and fourth stage larvae)

*Bunostomum phlebotomum* - (including third and fourth stage larvae)

##### And adult stages of:

*Trichuris* spp. - Whipworm

*Strongyloides papillosus* - Intestinal worm

*Nematodirus* spp. - Thin necked intestinal worm

#### ADULT LIVER FLUKE

*Fasciola hepatica*

#### LUNGWORMS

*Dictyocaulus viviparus* (adult and immature, including inhibited stages).

#### EYEWORM

*Thelazia* spp. (adult stages).

#### SUCKING LICE:

*Linognathus vituli* - Long nosed sucking louse

*Haematopinus eurysternus* - Short nosed sucking louse

*Solenopotes capillatus* - Little blue sucking louse.

#### MITES:

*Sarcoptes scabiei var bovis*

#### SCREW WORM FLY:

*Chrysomya bezziana* (parasitic larval stages. Not reported in Australia but present in Papua-New Guinea)

#### CATTLE TICK:

*Rhipicephalus (Boophilus) microplus* - Cattle tick

When used as directed for control of internal parasites of cattle, CATTLEMAX® PLUS INJECTION FOR CATTLE concurrently aids in control of:

#### MITES:

*Chorioptes bovis*

#### BITING LICE:

*Bovicola (Damalinia) bovis*

#### BEFORE USING THIS PRODUCT, READ ALL DIRECTIONS ON THE CARTON.

### DIRECTIONS FOR USE:

**RESTRAINT:** DO NOT USE in ewes, which are producing or may in the future produce.

The recommended dose is 1 mL CATTLEMAX® PLUS INJECTION FOR CATTLE per 50 kg liveweight (equivalent to 200 µg ivermectin and 2 mg clorsulon per kg liveweight) by subcutaneous injection.

This product is not to be used intravenously or intramuscularly.

Liveweight (kg)	Dose volume (mL)	Cattle Treated 500 mL	Liveweight (kg)	Dose volume (mL)	Cattle Treated 500 mL
Up to 100	2	250	351 - 400	8	62
101 - 150	3	166	401 - 450	9	55
151 - 200	4	125	451 - 500	10	50
201 - 250	5	100	501 - 550	11	45
251 - 300	6	83	551 - 600	12	41
301 - 350	7	71	601 - 650	13	38

Dose the mob according to the heaviest animal by liveweight in each group (cows, bulls, calves, heifers, etc). Do not underdose.

A representative sample of cattle should be weighed before treatment either with scales or with a weighband.

Cattle heavier than 650 kg should be dosed at 1 mL per 50 kg.

Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

**SUSTAINED ACTIVITY:** At the recommended dose rate, CATTLEMAX® PLUS INJECTION FOR CATTLE effectively controls infection with *Ostertagia* spp. and *Cooperia* spp. acquired up to 7 days after treatment, and *Dictyocaulus viviparus* and *Chrysomya bezziana* acquired up to 14 days after treatment.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER THE APPROPRIATE LEGISLATION**

#### WITHHOLDING PERIODS:

**MEAT:** DO NOT USE LESS than 28 days before slaughter for human consumption.

**MILK:** Zero (0) days

**TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 42 days before slaughter for export. Before using this product confirm the current ESI from either Loveland Agri Products Pty Ltd on 1800 448 892 or the APVMA website ([www.apvma.gov.au/residues](http://www.apvma.gov.au/residues)).**