

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



**Dalgety**  
Animal Health



# MOXIMAX<sup>®</sup>

LA INJECTABLE FOR SHEEP

**ACTIVE CONSTITUENTS: 20 g/L MOXIDECTIN**

For the treatment and control of roundworms, nasal bot and itchmite in sheep. For protection against severe challenge by *Haemonchus contortus* (barber's pole worm) for up to 4 months.

Resistance may develop to any chemical.

**WORM CONTROL**



10081186

[dalgetyah.com.au](http://dalgetyah.com.au)

**500 mL**

**SAFETY DIRECTIONS:**

Poisonous if swallowed. May irritate the nose and throat. Will irritate eyes and skin. Avoid contact with eyes and skin. Do not inhale. Wash hands after use.

**FIRST AID:** If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

**ADDITIONAL USER**

**SAFETY:** Additional information is listed in the Safety Data Sheet available from a retailer or the manufacturer.

**WARNING: AVOID SELF-INJECTION**  
Accidental self-injection may cause a persistent inflammatory or allergic response.

Medical advice should be sought as soon as possible on the management of all instances of self-injection, especially deep injections, those near a joint or those associated with bruising. The application of gentle pressure with absorbant

material, e.g. facial tissues, to the needle puncture area will swab up unabsorbed product. Strong squeezing of the site should be avoided. The damaged area should be thoroughly cleaned and a suitable antiseptic applied.

**ENVIRONMENT**

**PROTECTION:** Very toxic to aquatic life. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

**DISPOSAL:** If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Discarded needles/sharps should immediately be placed in a designated and appropriately labelled 'sharps' container. The container should be of a type to reduce the possibility of injury to handlers during collection and disposal. Incineration is the preferred method of disposal, otherwise sharps should be buried at a suitable site, such as an on-farm chemical disposal pit located away from water courses.

**STORAGE:** Store at or below 30 °C (room temperature). Protect from light.

**Loveland Agri  
Products Pty Ltd**  
Level 5, Building A, 26  
Talavera Road,  
Macquarie Park, NSW, 2113  
Tel: (02) 9889 5400  
Product Support  
Tel: 1800 44 88 92  
**dalgetyah.com.au**

APVMA Approval No.:  
88432/121304

[B]:  
Expiry:

**CAUTION**  
**KEEP OUT OF REACH OF CHILDREN**  
**FOR ANIMAL TREATMENT ONLY**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# MOXIMAX<sup>®</sup>

## LA INJECTABLE FOR SHEEP

**ACTIVE CONSTITUENTS: 20g/L MOXIDECTIN**

For the treatment and control of roundworms, nasal bot and itchmite in sheep.  
For protection against severe challenge by *Haemonchus contortus* (Barber's pole worm) for up to 4 months.

**INDICATIONS:** MOXIMAX<sup>®</sup> LA INJECTABLE FOR SHEEP contains moxidectin, a second generation of the macrocyclic lactone family, it is effective against sensitive strains of the following parasites and provides protection against severe challenge by *Haemonchus contortus* (barber's pole worm) for up to 4 months.

### INTERNAL PARASITES

Mature (adult) and Immature (L4)  
*Haemonchus contortus* (barber's pole worm)  
*Trichostrongylus* spp. (black scour worm)  
*Trichostrongylus axei* (stomach hair worm)  
*Trichostrongylus colubriformis*  
*Teladorsagia (Ostertagia) spp.* (small brown stomach worm)  
*Teladorsagia (Ostertagia) circumcincta*  
*Ostertagia lyrata*  
*Cooperia* spp. (small intestinal worm)  
*Cooperia oncophora*  
*Nematodirus* spp. (thin necked intestinal worm)  
*Nematodirus filicollis*  
*Nematodirus spathiger*  
*Strongyloides papillosus* (intestinal threadworm)  
*Oesophagostomum columbianum* (nodule worm)  
*Chabertia ovina* (large mouthed bowel worm)  
*Dictyocaulus filaria* (large lungworm)

### ADULT NEMATODES

*Trichostrongylus rugatus*  
*Trichostrongylus vitrinus*  
*Ostertagia trifurcata*  
*Cooperia curticei*  
*Nematodirus abnormalis*  
*Nematodirus helvetianus*  
*Oesophagostomum venulosum* (large bowel worm)  
*Trichuris ovis*

### EXTERNAL PARASITES

*Psorergates ovis* (itchmite)  
*Oestrus ovis* (nasal bot)

**PROTECTION PERIOD:** When MOXIMAX® LA INJECTABLE FOR SHEEP is used at the recommended dose rate as a single subcutaneous injection, it prevents reinfection of sheep with nematodes as in the following table:

Protection	<i>H. contortus</i>	<i>T.(O.) circumcincta</i>	<i>T. colubriformis</i>
Prevents reinfection of sheep with worms	Not less than 91 days	Not less than 91 days	Up to 49 days

MOXIMAX® LA INJECTABLE FOR SHEEP has been shown to protect sheep against severe challenge by *H. contortus* for up to 4 months.

MOXIMAX® LA INJECTABLE FOR SHEEP is effective against benzimidazole and levamisole resistant strains of *H. contortus*, *T.colubriformis* and *T.(O.) circumcincta*.

MOXIMAX® LA INJECTABLE FOR SHEEP has shown efficacy against a number of avermectin resistant strains of *H. contortus* and *Teladorsagia (Ostertagia)* spp. However, as MOXIMAX® LA Injectable for Sheep shares a similar mechanism of action to avermectins, it SHOULD NOT be used where avermectin resistance is present, except where specifically recommended by an advisor.

**RESISTANCE AND WORM CONTROL PROGRAMS:** It is advisable that a drench resistance test be conducted before any anthelmintic treatment is used.

Resistance may develop to any anthelmintic treatment. Ask your local veterinary practitioner or animal health advisor for recommended parasite management practices for your area.

**RESTRAINT:** DO NOT USE in ewes, which are producing or may in the future produce milk that may be used or processed for human consumption.

**RETREATMENT INTERVAL:** Must not be retreated for at least 91 days after administration.

**CONTRAINDICATION:** MOXIMAX® LA Injectable for Sheep is not recommended for use in goats as safety and efficacy have not been evaluated.


**DOSAGE AND ADMINISTRATION:** Use the contents within 24 hours of first broaching the vial. Discard the unused portion.

**DOSE RATE:** 1 mg moxidectin/kg liveweight (1 mL/20 kg liveweight).

MOXIMAX® LA Injectable for Sheep must be administered by subcutaneous (under the skin) injection at the base of the ear or high on the neck behind the ear.

The product is ready to use. Administer the dose according to the dosage table using the special MOXIMAX® LA Injectable for Sheep Injector gun or similar Moxidectin Long Acting · Injector gun. Do not use any other injector. Check dose rate and equipment before treatment commences.

Sheep should be weighed prior to dosing and treated according to the weight range bracket in the dosage table below. Do not underdose.



**DOSAGE TABLE**

Weight range (kg)	Dose volume (mL)	Doses per pack 500 mL
*20	1	500
21-30	1.5	333
31-40	2	250
41-50	2.5	200
51-60	3	166
61-70	3.5	142
71-80	4	125
81-90	4.5	111

**\*DO NOT USE IN LAMBS LESS THAN 20 kg.**

**Animals heavier than 90 kg to be dosed at 0.5 mL per 10 kg.**

MOXIMAX® LA INJECTABLE FOR SHEEP has a wide margin of safety when used as recommended.

MOXIMAX® LA INJECTABLE FOR SHEEP is safe to use concurrently with other treatments including mineral supplements and vaccines. It is recommended that vaccines are injected on the opposite side of the neck of MOXIMAX® LA Injectable for Sheep.

**CAUTION TO AVOID CARCASS DAMAGE:**

- Rinse all injection syringes, ensure needles and draw off tubes are free of dirt and unused product before and after use;
- Sanitise plastic injection apparatus by immersing in 500 mL of water plus 20 mL household bleach (4%w/v available chlorine) for 1 hour. Flush with cool boiled water before use.
- Maintain cleanliness at all times.
- Keep needles sharp and clean, replace frequently.
- Avoid injection of animals during wet weather or under dusty conditions as far as possible.
- Inject the product under the skin, NOT into the muscle tissue.
- Inject at base of the ear or high on the neck behind the ear.

**WITHHOLDING PERIODS:**

**MEAT: DO NOT USE less than 91 days before slaughter for human consumption.**

**MILK: DO NOT USE in ewes, which are producing or may in the future produce milk that may be used or processed for human consumption.**

**TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 133 days before slaughter for export.**

**Before using this product, confirm the current ESI from the manufacturer on 1800 448 892 or the APVMA website ([www.apvma.gov.au/residues](http://www.apvma.gov.au/residues))**

**SAFETY DIRECTIONS:** Poisonous if swallowed. May irritate the nose and throat. Will irritate eyes and skin. Avoid contact with eyes and skin. Do not inhale. Wash hands after use.

**FIRST AID:** If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

**ADDITIONAL SAFETY INFORMATION:** Avoid self-injection. If self-injection occurs, seek medical advice as soon as possible. Additional information is listed in the Safety Data Sheet available from a retailer or the manufacturer.

**WARNING: AVOID SELF-INJECTION.** Accidental self-injection may cause a persistent inflammatory or allergic response.

Medical advice should be sought as soon as possible on the management of all instances of self-injection, especially deep injections, those near a joint or those associated with bruising. The application of gentle pressure with absorbant material, e.g. facial tissues, to the needle puncture area will swab up unabsorbed product. Strong squeezing of the site should be avoided. The damaged area should be thoroughly cleaned and a suitable antiseptic applied.

**ENVIRONMENTAL PROTECTION:** Very toxic to aquatic life. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

**DISPOSAL:** If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

Discarded needles/sharps should immediately be placed in a designated and appropriately labelled 'sharps' container. The container should be of a type to reduce the possibility of injury to handlers during collection and disposal. Incineration is the preferred method of disposal, otherwise sharps should be buried at a suitable site, such as an on-farm chemical disposal pit located away from water courses.

**STORAGE:** Store at or below 30°C (room temperature). Protect from light.

**CONDITIONS OF SALE:** This product should only be used in accordance with the label and manufacturer's instructions. Product performance may be affected by factors which are beyond the control of Loveland Agri Products Pty Ltd and its Associates (as that term is defined in the Corporations Act 2001 Cth) (Associates). If symptoms persist, please consult your veterinarian or animal health specialist. In no case shall Loveland Agri Products Pty Ltd or its Associates be liable for consequential, incidental, special or exemplary damages or losses. All sales (wholesale and retail) are governed by the General Terms and Conditions of Sale available at [lovelandagriproducts.com.au](http://lovelandagriproducts.com.au) or from your retailer upon request. These terms can only be varied by Loveland Agri Products Pty Ltd.

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: May cause damage to organs through prolonged or repeated exposure; Do not eat, drink or smoke when using this product; Use personal protective equipment as required; If swallowed rinse mouth and DO NOT induce vomiting; If skin irritation occurs or eye irritation persists get medical advice; Not combustible - use extinguishing media suited to burning materials; Store in a dry, well ventilated place and keep cool.

APVMA Approval No.: 88432/121304

B:

Exp:

**Loveland Agri Products Pty Ltd**

Level 5, Building A, 26 Talavera Road, Macquarie Park, NSW, 2113

Tel: (02) 9889 5400

Product Support Tel: 1800 44 88 92

**[dalgetyah.com.au](http://dalgetyah.com.au)**

IN A TRANSPORT EMERGENCY

**DIAL 000**

POLICE OR FIRE BRIGADE

FOR SPECIALIST ADVICE IN AN EMERGENCY

**1800 033 111**

24 HOURS AUSTRALIA WIDE



Please note instruction leaflet  
and carton only are recyclable.